

CAT Nº: 6036

FORMULA PER VIAL

Cetrimide	5 mg
Fucidin	5 mg
Cephaloridine	25 mg

PREPARATION

Aseptically reconstitute 1 vial with 5 ml of warm sterile distilled water. Mix gently until complete dissolution and aseptically add to 500 ml of PSEUDOMONAS AGAR BASE (Cat. 1356). Mix well and distribute into sterile containers.

PRODUCT	CAT.	PACK SIZE
C.F.C SUPPLEMENT	6036	10 vials
PSEUDOMONAS AGAR BASE	1356	500 g

MICROBIOLOGICAL TEST

The following results were obtained from type cultures in the performance of PSEUDOMONAS AGAR BASE, with the supplement added, after incubation at a temperature of 25± 1°C and observed after 44±4 hours.

Microorganisms	Growth	Recovery
<i>Escherichia coli</i> ATCC 25922	Inhibited	-
<i>Pseudomonas aeruginosa</i> ATCC 27853	Good	≥ 70%
<i>Pseudomonas aeruginosa</i> ATCC 25819	Good	≥ 70%
<i>Pseudomonas aeruginosa</i> ATCC 25608	Good	≥ 70%

BIBLIOGRAPHY

Goto, S., and S. Enomoto. 1970. Nalidixic Acid Cetrimide Agar. A New Selective Plating Medium for the Selective Isolation of *Pseudomonas aeruginosa*. Japan. J. Microbiol. 14: 65 - 72.



STORAGE

When stored as indicated, supplements remain stable until the stated expiry date shown on the label.

